




Available online at  
 ScienceDirect  
www.sciencedirect.com

Elsevier Masson France  
  
www.em-consulte.com



## GUIDELINES

# Consensus of the French Society of Gerontology and Geriatrics and the French Society of Cardiology for the management of coronary artery disease in older adults

Consensus d'experts de la Société française de gériatrie et gérontologie (SFGG) et de la Société française de cardiologie (SFC) sur la prise en charge de la maladie coronaire chez le sujet âgé

Olivier Hanon<sup>a,\*</sup>, Cécile Baixas<sup>b</sup>, Patrick Friocourt<sup>a</sup>,  
Didier Carrié<sup>b</sup>, Jean-Paul Emeriau<sup>a</sup>, Michel Galinier<sup>b</sup>,  
Joel Belmin<sup>a</sup>, Pascal de Groote<sup>b</sup>, Athanase Bénétos<sup>a</sup>,  
Patrick Jourdain<sup>b</sup>, Gilles Berrut<sup>a</sup>,  
Jean-François Aupetit<sup>b</sup>, Guillaume Jondeau<sup>b</sup>,  
Nicolas Danchin<sup>b</sup>, Françoise Forette<sup>a</sup>,  
Michel Komajda<sup>b,\*\*</sup>

<sup>a</sup> French Society of Gerontology and Geriatrics, France

<sup>b</sup> French Society of Cardiology, France

Received 4 September 2009; accepted 4 September 2009

## KEYWORDS

Geriatrics;

**Summary** Coronary heart disease is a common and serious condition in patients aged over 80 years. The presenting clinical symptoms are all the more atypical and the prognosis poorer when it occurs in patients with multiple comorbid diseases. The presence of comorbidities dictates the need for a standardized geriatric assessment to screen for the existence of underlying

**Abbreviations:** ACS, acute coronary syndrome; ADL, activities of daily living; ARA, angiotensin II receptor antagonist; ASSENT-3, Assessment of the Safety and Efficacy of a New Treatment Strategy with Percutaneous Coronary Intervention; CURE, Clopidogrel in Unstable angina to prevent Recurrent Events; GDS, geriatric depression scale; ECG, electrocardiogram; GISSI-3, Third Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico; HYVET, hypertension in the very elderly trial; IADL, instrumental activities of daily living; IONA, economic evaluation of the impact of nicorandil in angina; LMWH, low-molecular-weight heparin; MMSE, Mini Mental Status Examination; MNA, Mini Nutritional Assessment; PROSPER, pravastatin in elderly individuals at risk of vascular disease; PTCA, percutaneous transluminal coronary angioplasty; SCGA, standardized comprehensive geriatric assessment; SHOCK, Should we emergently revascularize Occluded coronaries for Cardiogenic shock?.

\* Corresponding author at: Department of Geriatrics, université Paris-Descartes, hôpital Broca, 54-56, rue Pascal, 75013 Paris, France.

\*\* Corresponding author at: Department of Cardiology, Pitié Salpêtrière Hospital group, 47–83, boulevard de l'Hôpital, 75651 Paris cedex 13, France.

E-mail addresses: [olivier.hanon@brc.ap-hop-paris.fr](mailto:olivier.hanon@brc.ap-hop-paris.fr) (O. Hanon), [michel.komajda@psl.aphp.fr](mailto:michel.komajda@psl.aphp.fr) (M. Komajda).

Coronary artery disease;  
Myocardial infarction;  
Aging;  
Elderly

frailty. The available scientific data were obtained during studies that included few subjects aged over 80 years. These recommendations are therefore mainly extrapolated from results obtained in younger populations. The pharmacological management and revascularization strategy for coronary heart disease in octogenarians is basically the same as in younger subjects. Epidemiological studies all concur that available therapies are underutilized despite the fact that this population has a high cardiovascular risk. Specific precautions for use must be respected because of the comorbidities and age-related changes in pharmacokinetics or pharmacodynamics. Generally, the therapeutic strategy in coronary heart disease is based not on the patient's real age, but rather on an individual analysis taking into account the severity of the coronary disease, comorbidities, the risk of drug misadventures, patient life expectancy and quality of life.

## Introduction

Mortality due to ischaemic heart disease fell by 16% in France between 1988–1990 and 1995–1997, but this condition still causes 27% of deaths from cardiovascular disease. This mortality increases with age and is always higher in men [1]. Nearly one-third [2–4] of patients admitted for myocardial infarction are over 75 years old and represent nearly 50% of hospital mortality due to this disease. Patients over 85 years have six times more cardiogenic shocks, three times more strokes, twice as many major bleeding events and three to nine times higher mortality [5,6].

Few data are available in the literature about coronary insufficiency in adults aged over 80 years. Old age is an exclusion criterion in most randomized controlled trials on ACS. This article, compiled from a review of the literature, is an 'expert consensus' of the French Societies of cardiology and geriatrics/gerontology on the specific features of management of coronary heart disease in patients aged over 80.

## Clinical approach

The presentation of coronary insufficiency in the older subject often differs from that in younger adults and atypical clinical features are often observed.

### Angina pain

Angina pain remains the most frequent inaugural sign of coronary insufficiency in the elderly, although it has a lower incidence than in younger adults. It may be more difficult to analyse because of communication difficulties and in particular the presence of cognitive impairment, which may cause patients to 'forget' symptoms. Atypical pathological features are frequent in older adults, particularly in women: gastrointestinal disorders, asthenia and deterioration in general health status may be the main presenting symptoms. Ischaemic heart disease may also lead to block-pnea, acute pulmonary oedema, rhythm disorders or even sudden death [7].

### Silent ischaemia

Silent ischaemia is more frequent in the elderly [8] because of impaired sensory nerve function, deterioration in cortical function or dysautonomia. Coronary heart disease may

be asymptomatic when physical activity is reduced below the ischaemic threshold by a sedentary lifestyle or locomotor disability. The incidence of confounding diseases [9] (spondylosis or shoulder osteoarthritis, chest wall pain or gastrointestinal diseases) makes clinical analysis more difficult.

### Myocardial infarction

Diagnosis of myocardial infarction may also be difficult in older adults and the presenting symptom may be a complication (acute pulmonary oedema, stroke, acute ischaemia of the lower limbs, heart rhythm disorder or sudden death) or a gastrointestinal or neurological symptom (confusional syndrome, behavioural disorders, syncope or vertigo or even simple headaches [10]). Atypical symptoms often lead to delays in treatment [11].

### Clinical examination

The clinical examination determines the distribution of the atheromatous disease (arterial murmur, detection of abdominal aortic aneurysm). The physical examination may also reveal arrhythmia or signs of heart failure. A measurement of seated and then standing blood pressure is indicated (to test for orthostatic hypotension, defined as a fall of 20 mmHg in systolic blood pressure and/or 10 mmHg in diastolic blood pressure). Finally, a precise neurological examination should be performed. This should be completed by a SCGA.

## Geriatric approach

The management of older adults with coronary insufficiency has different objectives depending on the patient. In autonomous elderly subjects with no severe concomitant disease, it is identical to that of younger adults. The aim of management of a dependent older adult with multiple comorbid diseases is to preserve vital functions while avoiding adverse drug reactions.

### Global assessment

This essential assessment is often difficult to perform in an emergency setting and must be completed when the patient's condition has stabilized, where necessary by a specialized team. It assesses medical and psychosocial items and uses simple tests for the rapid detection of concomitant

diseases, an evaluation of the patient's autonomy and his/her social conditions. The SCGA makes it possible to evaluate frailty characterized by a reduction in physiological capacities to adapt to stress or changes in the environment associated or not with organ failure. The main items of the SCGA are the following.

### Evaluation of cognition

The MMSE (Folstein's MMSE [12]), scored out of 30, constitutes a simple, rapid and standardized test for detecting cognitive impairment (Appendix 1) [13]. The threshold value depends on the patient's age and sociocultural level. A score over 27 is considered to be normal. A score below 24 is abnormal and dictates the need for a detailed assessment of cognitive function in a specialized unit. A score of between 24 and 27 should be considered abnormal for patients who are not extremely old (< 80 years), with a high sociocultural level and who have disturbances of memory or other cognitive functions. An abnormal MMSE score shows cognitive dysfunction not necessarily related to dementia. An abnormal score therefore calls for specialized evaluation and is not sufficient in itself to establish diagnosis.

### Evaluation of dependence

Evaluation of dependence specifies limitations that may hinder complementary investigations, worsen prognosis [14] and impact the therapeutic approach. Dependence may be assessed with Activities of Daily Living scales (IADL or ADL) by interviewing the patient and his/her close contacts. There is a shortened four-item form of the IADL [15] based on: use of the telephone, means of transport, medication use and management of money (Appendix 2). The ADL scale provides information about personal hygiene, dressing, use of toilets, transferring, urine and bowel continence and eating. The subject is considered to be dependent as soon as he/she needs human help to carry out the activity concerned (Appendix 3) [16].

### Somatic geriatric evaluation

This takes into account the evaluation of gait disorders and the risk of falling (unipedal stance test to evaluate the capacity to stand on one leg for 5 seconds). Likewise, nutritional status is assessed using validated scales (MNA scale) for the assay of plasma albumin or prealbumin [17]. Body weight must be interpreted after taking into account possible malnutrition and any fluid retention or dehydration.

### Psychological evaluation

Depression is very frequent in older adults and may impact the manifestations of coronary disease and its treatment. It represents an independent risk factor for cardiovascular mortality [18]. Its detection and management are therefore essential in older adults. The simplified GDS provides an initial assessment (Appendix 4).

### Evaluation of life circumstances

The purpose of this evaluation is to:

- estimate adherence to the proposed treatment. If the patient cannot manage his/her own treatment, it must be

determined how (preparation, use of a tablet container) and by whom (family, domestic assistant, carer, nurse) it will be administered;

- specify the procedures for preparing food and meals;
- evaluate the patient's state of isolation and his/her access to different care services.

### Community health management

It is important to evaluate the patient's social condition and take into account the role of caregivers (family relations or friends). This management involves informing and educating the patient's close contacts about the disease to make sure that he/she takes the medication and permit rapid detection of any change in symptoms and thresholds for the triggering of symptoms. This may require implementation of a home-help system. Different types of care and support are available including help to ensure personal hygiene from a nurse or nursing auxiliary, supply of medications (nurse), meals-on-wheels, domestic help (home helper) or care-giving (carer).

Depending on the SCGA data, the patient may be seen by a geriatrician in order to organize management and appropriate follow-up. Such management of the frail elderly has been shown to result in a significant reduction in mortality and admission to institutions [19].

### Additional diagnostic and prognostic investigations

#### Electrocardiogram (ECG)

The ECG is rarely normal at baseline in older persons with coronary disease. Interpretation of repolarization is often hindered by the presence of left ventricular hypertrophy, electrical sequelae of infarction, a left bundle branch block, electrosystolic pacing, treatment by digitalis preparations or electrolyte disorders. The resting ECG may, however, be normal [20]. In this case, repeated recordings at short intervals and clinical and laboratory examinations should improve diagnosis. The ECG also provides prognostic information. For instance, the risk of coronary events is raised in the case of ST-segment depression greater than 1 mm [21].

#### Exercise ECG

An exercise ECG may rarely be performed in older adults who often have mobility disorders, joint pain, physical deconditioning and concomitant neurological, vascular or respiratory disorders [22]. No study of the sensitivity and specificity of the exercise test has been performed in the very elderly. Those evaluating its prognostic value gave contradictory results [23,24].

#### Myocardial perfusion scintigraphy

Myocardial perfusion scintigraphy, at rest or after dipyridamole or dobutamine stress, has a good sensitivity and specificity in subjects aged over 80 years for the detection of significant coronary stenosis [25] and the evaluation of the prognosis [26], although this examination is not always available.

## Stress echocardiography

Stress echocardiography may also be proposed in the elderly. Physical exercise may be used as stress factor although pharmacological stress testing is more often performed because of functional limitations. The sensitivity and specificity of exertional echocardiography are similar to those in populations of younger patients [27]. After 75 years, dobutamine stress echocardiography is accompanied by an increased incidence of hypotension and ventricular arrhythmia [28].

## Other evaluation procedures

The role of coronary computed tomography angiography and magnetic resonance imaging in the diagnosis of the coronary insufficiency is undergoing evaluation in older adults in particular, because of the renal risks related to iodine injection required for the computed tomography scan.

## Other complementary examinations

Assays of blood glucose levels and a lipid assessment are useful for the evaluation of cardiovascular risk. Serum creatinine concentrations and creatinine clearance calculated with the Cockcroft-Gault formula are used to assess renal function. Impaired renal function, in particular when severe (clearance < 30 mL/min), is a factor of poor prognosis, increasing the risk of adverse drug events especially if coronary angiography is performed. Rehydration is essential before conducting this examination, as soon as the clearance is less than 60 mL/min.

## Markers of ischaemia and myocardial necrosis

Assay of biomarkers of myocardial ischaemia (creatinine phosphokinase, troponin) contribute to the diagnosis and prognosis for patients with ACS. The procedures for using and interpreting the results of these assays are not affected by age, although the presence of macro creatine kinases, leading to false positives, is more frequent in elderly women [29]. In the case of severe renal insufficiency, a slight rise in troponin T may occur outside any ACS [30].

## Coronary angiography

Coronary angiography is the reference diagnostic examination for the detection of coronary lesions. It has a higher morbidity and mortality rate in octogenarians. The indication for coronary angiography must therefore be established after evaluating the patient's conditions, mainly with respect to possible revascularization.

## Medical treatment of ACS in the older subject

### Definition

ACS [31] in older adults comprises:

- ST-segment elevation ACS: prolonged suggestive chest pain (> 20 minutes' duration) accompanied by persistent ST-segment elevation (or new left bundle branch block);

increase in cardiac biomarkers (in particular troponin I or T) [32];

- non-ST-segment elevation ACS: prolonged suggestive chest pain or chest pain of recent onset (de novo angina) or worsening of angina generally accompanied by electrical modifications without persistent ST-segment elevation; increase in cardiac biomarkers (in particular troponin I or T) [33,34].

## Lack of 'evidence-based' data

Older adults are under-represented in the main studies in ACS. More than half of the trials conducted between 1996 and 2000 included no patient aged over 75 years [28]. It is not therefore known if the algorithms and guidelines established with cohorts of younger patients are applicable to older subjects. Likewise, the best endpoint for use in this population with a reduced life expectancy is not known (mortality, cardiovascular events or quality of life?).

## Medical management

The different epidemiological studies all underline the delayed management and under-utilization of medication in ACS. Compared with younger patients, those over 75 years are less often given beta-blockers, aspirin, clopidogrel, statins, thrombolysis (in the case of ST-segment elevation ACS) and glycoprotein IIb/IIIa antagonists [2–4]. Conversely, elderly subjects often receive more calcium antagonists and angiotensin-converting enzyme inhibitors [4], probably because of the increased prevalence of hypertension and heart failure. Likewise, increased use of LMWHs is observed despite the increased bleeding risk in this population.

The optimum medical management of older adults with ACS is not very different from that of younger subjects. However, the frailty of this population dictates the need for increased rigor in respecting contraindications, careful institution of the different therapeutic classes and monitoring for possible side-effects.

## Platelet-aggregation inhibitors

### Aspirin

The prescription of aspirin is recommended [35,36] from the acute phase, at a dosage of between 75 and 325 mg/day in the absence of allergy, exacerbation of peptic ulcer or bleeding disorders, and must be continued over the long term. Its side-effects (mainly gastrointestinal) are uncommon at low doses (75–150 mg/day) but must be carefully sought.

### Clopidogrel

The prescription of clopidogrel monotherapy is recommended in the case of contraindications to aspirin. The combination of clopidogrel with low doses of aspirin has not been evaluated in subjects over 80 years. Taking into account the divergent results of the CURE [37] and PCI-CURE studies [38] in patients aged over 65 years, dual antiplatelet therapy may only be recommended on an individual basis after taking into account the benefit-to-risk ratio for each patient.



The optimum duration of treatment with the aspirin–clopidogrel combination has not been well evaluated in older adults, and must take into account the patient's general status and/or the implantation of a drug-eluting stent. There are no data evaluating the benefit and safety of loading doses in patients over 80 years.

### Withdrawal of antiplatelet therapy in older adults

This is frequently necessary because of the multiple comorbid diseases in this population. Discontinuation of platelet aggregation inhibitors is a major risk factor for thrombosis of all coronary stents and in particular late thrombosis in drug-eluting stents. Recommendations exist about the management of oral antiplatelet therapy in patients with coronary stents, in particular in patients requiring surgery, although these do not specifically concern elderly patients [39].

### Glycoprotein IIb/IIIa antagonists

Few randomized studies have evaluated the benefit of glycoprotein IIb/IIIa antagonists in patients over 75 years. In patients with ST-segment elevation ACS, they may be used in primary angioplasty with coronary stenting [29,40] after careful analysis of the trade-off between benefit and risks, and ruling out contraindications (stroke, surgery or recent trauma, coagulation disorders, hepatic insufficiency, active haemorrhage, severe arterial hypertension). On the contrary, their prescription, in combination with fibrinolytics, is clearly contraindicated in patients over 75 years [29].

Subjects over 75 years, admitted for non-ST-segment elevation ACS also seem to benefit from adjunctive anti-glycoprotein IIb/IIIa therapy with early revascularization at the price, however, of increased bleeding [41].

## Anticoagulants

### Unfractionated heparin

Unfractionated heparin is recommended in ST-segment elevation ACS whatever revascularization procedure is used, provided there are no immunoallergic or haemorrhagic contraindications [42]. In subjects aged over 75 years with ST-segment elevation ACS undergoing treatment with fibrinolysis, unfractionated heparin was better tolerated than the LMWH enoxaparin in terms of the risk of intracranial haemorrhage [37]. In these high-risk patients, clinical and laboratory monitoring should be reinforced for signs of visceral or cerebral bleeding.

### Low molecular weight heparin

LMWHs, in particular enoxaparin, are recommended in young patients with ACS [43]. After the age of 75 years, in combination with thrombolysis during the acute phase of myocardial infarction, a subgroup of the ASSENT-3 study suggested that LMWH and unfractionated heparin had similar benefits in terms of the composite endpoint of death, in-hospital reinfarction, refractory ischaemia, intracranial haemorrhage and major bleeding [44]. However, the risk of intracranial haemorrhage was higher with enoxaparin than with unfractionated heparin in older adults in ASSENT-3 Plus [42].

The advantages of LMWH are their ease of use and the lower risk of heparin-induced thrombocytopenia compared with unfractionated heparin. Conversely, they must be prescribed with caution in patients aged over 75 years because of their renal elimination. The dosage must be adjusted according to renal function and anti-Xa activity to avoid accumulation responsible for haemorrhagic accidents [45]. A creatinine clearance less than 30 mL/min is an absolute contraindication for curative doses whereas a clearance of between 30 and 60 mL/min is a relative contraindication (with obligatory control of anti-Xa activity). The duration of anticoagulant treatment ranged from 48 hours to eight days depending on the studies and the type of revascularization procedure which is not codified in the elderly.

### Direct antithrombotics and factor Xa inhibitors

Although they are not currently recommended in elderly patients with ACS [36], direct thrombin inhibitors appear promising [46,47]: fondaparinux appears to have a similar efficacy to that of enoxaparin in patients with non-ST-segment elevation ACS, with a lower bleeding risk, particularly in older subjects [48]. In ST-segment elevation ACS, although safety was the same, the benefit was not as marked as in the subgroup of patients undergoing primary angioplasty [49].

### Vitamin K antagonists

Vitamin K antagonists are not part of the treatment armamentarium in ACS. They must be stopped where necessary and relayed by heparin anticoagulation.

### Beta-blockers

Irrespective of the type of ACS (ST-segment elevation or non-ST-segment elevation), beta-blockers have clear short- and long-term benefits in terms of reducing recurrence of myocardial infarction and increasing survival. Although few studies have included patients over 75 years, the data obtained from meta-analyses and registries show a benefit whatever the patient's age. They are recommended for first-line treatment [29], in particular in the case of left ventricular dysfunction [50] after ruling out all the usual contraindications, such as decompensated heart failure, asthma and advanced chronic obstructive pulmonary disease, and severe conduction disorders, which must be sought with particular care in the elderly.

Close monitoring of blood pressure, heart rate and the ECG are necessary in this setting. Monitoring for the onset of bradycardia is necessary in the case of concomitant prescription of acetylcholinesterase inhibitors. Institution of beta-blockers must be delayed if there are overt signs of water and sodium retention. Treatment should be initiated cautiously with gradual up-titration of the dose to a target resting heart rate of approximately 60 beats per minute.

## Inhibitors of the renin-angiotensin system

### Angiotensin-converting enzyme inhibitors

Angiotensin-converting enzyme (ACE) inhibitors are recommended during the first 24 hours of onset of ACS if there are clinical signs of heart failure and/or a deterioration

in left ventricular systolic function (left ventricular ejection fraction <40%) and/or anterior-wall infarction [51,52]. They are indicated after myocardial infarction in all patients and indefinitely, in the absence of contraindications. There are few data specific to patients aged over 75 years and these concern subgroup analysis [53]. ACE inhibitors should be initiated cautiously in older adults and the dose gradually increased according to renal function. In general, the daily dosage should be reduced by half compared with that used in younger patients. Treatment should be initiated long after any episode of sodium depletion, starting with a low dosage, which is then gradually up-titrated to the maximum tolerated dose. At the same time, the continuation or interruption of other therapies such as diuretics, calcium antagonists and nitrates should be re-evaluated. Prescription of non-steroidal anti-inflammatory drugs is contraindicated because of the risk of renal insufficiency and hyperkalaemia. Blood pressure (seated and standing), renal function and blood electrolytes must be monitored carefully and very regularly during the first months after myocardial infarction, and systematically in the case of acute intercurrent episodes of fever, diarrhoea, dehydration, vomiting, etc.

### Angiotensin II receptor antagonists (ARAs)

Two trials [54,55] evaluated the role of ARAs in post-infarction heart failure, and included a significant number of patients aged over 75 years. The results showed that ARAs had a similar efficacy to ACE inhibitors irrespective of the patient's age. Their use is therefore restricted to post-infarct patients intolerant to ACE inhibitors and younger subjects. They must be instituted at a low dose. Upward dose titration and monitoring procedures are similar to those recommended for ACE inhibitors.

### Aldosterone antagonists

No significant differences in benefit were observed between different age groups when eplerenone was used post-infarction in patients with signs of heart failure or left ventricular dysfunction. The chosen threshold was, however, particularly low (65 years) [56]. Because of the risk of hyperkalaemia due to renal function impairment in the elderly, the trade-off between risks and benefit should be carefully assessed and the contraindications respected in the absence of any specific study.

### Statins

Few scientific data are available showing the benefit of statins administered to ACS patients aged over 75 years. However, subgroup studies suggest that their beneficial effect is at least the same as that obtained in younger patients [57]. The target low-density lipoprotein cholesterol value and the doses required to reach it have not been determined.

### Calcium antagonists

Dihydropyridines have not been shown to have any benefit in patients with an ACS. They should only be prescribed within

the scope of their recognised indications, such as hypertension. They are not therefore recommended except in the specific case of confirmed coronary spasm.

The use of rate-limiting calcium antagonists (diltiazem, verapamil) may be considered when beta-blockers are contraindicated, with strict monitoring of the heart rate, ECG and blood pressure and for the possible onset of signs of heart failure.

### Nitrates

Nitrates have not been shown to improve outcomes in patients with ACS [58,59]. However, in the subgroup of patients aged over 70 years included in GISSI-3, nitrates administered within 24 hours of the onset of myocardial infarction significantly reduced cardiovascular complications [59].

They may be particularly useful in subgroups of ACS patients with recurrent or persistent myocardial ischaemia and/or left ventricular insufficiency and/or exacerbations of their high blood pressure [5]. They may be used temporarily at low doses to relieve pain, in the absence of hypotension or right ventricular infarction and with close monitoring of blood pressure and for the onset of headaches.

### Potassium channel agonists

No data are available in older adults with ACS. This treatment may, however, be considered in the case of clinically persistent angina if blood pressure values allow after the institution of the other pharmacological classes.

### Myocardial revascularization

Coronary reperfusion for myocardial infarction in older adults remains a controversial subject, mainly because of the under-representation or even exclusion of patients aged over 75 years in large clinical trials [60–62].

### Intravenous thrombolysis in myocardial infarction in older adults

Several studies and registries have underlined the increased risk of haemorrhage with advancing age during the hospital phase [63,64]. Nevertheless, thrombolysis significantly reduces 1-year mortality in patients over 75 years of age [65,66].

The risk of cerebral haemorrhage is twice as high after the age of 85 years but remains acceptable (approximately 3%) [67]. Factors predisposing to bleeding are: age, presence of comorbidities, malnutrition, severe arterial hypertension and brain trauma.

### Coronary angioplasty

The results of a recent meta-analysis of randomized trials [68] show that angioplasty in patients aged over 80 years has a greater benefit than thrombolysis, with, in particular, a better survival and a lower risk of stroke. These conclusions are coherent with those of observational studies [69–72]. This management strategy avoids the risks of haemorrhage due to thrombolysis. However, in older adults, angioplasty

is accompanied by a risk of more severe complications than in younger individuals (deaths, renal insufficiency, vascular complications) [73]. An evaluation of renal function before and after angioplasty is essential. It is particularly important to apply preventive measures with adequate hydration in the case of renal insufficiency. The indications for angioplasty during the acute phase in older adults must therefore be taken into account:

- clinical severity (recurrence of pain, heart failure);
- severity of the ischaemia (anterior territory or extension to the right ventricle, large rise in troponin);
- delay after the onset of signs (< 6 hours);
- physiological status (life expectancy, quality of life);
- presence of comorbidities (renal insufficiency, cognitive disorders, dependence, etc.).

### Coronary angioplasty with or without stent implantation

No randomized study has compared angioplasty with or without stenting in the very elderly subject with myocardial infarction.

### Revascularization in the case of cardiogenic shock

According to the SHOCK study, irrespective of the revascularization strategy (angioplasty or bypass) a reduction in mortality at six months is observed only in those aged under 75 years [74]. However, certain studies suggest a benefit on mortality after early revascularization in elderly patients selected according to individual criteria [75–77]. Revascularization, in this setting, can only be decided at the end of an individual decision process.

### Coronary artery bypass grafting

The perioperative morbidity and mortality of bypass grafting increase in older adults more especially if they have unstable angina or require emergency surgery [78,79].

### Summary

Angioplasty may be considered to be the technique of choice for emergency reperfusion. However, if this technique cannot be used, thrombolysis has benefits that should entail its wider use when the patient's physiological status permits.

## Treatment of chronic coronary disease

Medication for chronic ischaemic heart disease in older adults comprises two aspects:

- symptomatic treatments to reduce the frequency of angina pain and improve tolerance to exercise: nitrates, beta-blockers, calcium channel blockers, molsidomine, nicorandil, trimetazidine and ivabradine;
- secondary preventive treatments to improve outcomes by reducing morbidity and mortality: beta-blockers, aspirin, clopidogrel, statins, ACE inhibitors and ARAs.

### Treatment of angina attacks

For exertional angina, sublingual nitrates remain the basic treatment to relieve pain. The risk of orthostatic

hypotension after nitrate administration is increased in older adults, especially in the presence of aortic stenosis. They must be taken while seated. The elderly subject must receive advice about the factors precipitating angina attacks, how to take glyceryl trinitrate and the recommended treatment for intractable pain. The patient and his/her close relations must understand this information.

### Symptomatic treatment of stable angina pectoris

Few data are available in the literature about the antiangina efficacy of the different therapeutic classes in older adults. The conclusions of trials in the general population have therefore been extrapolated to the elderly [80].

### Beta-blockers

The use of beta-blockers is recommended in the absence of contraindications. Beta-blockers reduce the frequency of angina attacks and episodes of silent ischaemia [81,82]. Their dosage must be adjusted to maintain a resting heart rate of about 60 beats per minute and reduce the acceleration of heart rate during exercise [20,83]. Their administration should respect the precautions for use described in the section on ACS. A beta-1 selective beta-blocker should be preferred in patients with chronic and stable lower-limb arterial disease or diabetes. It is recommended not to suddenly discontinue beta-blocker treatment in patients with angina because of the increased risk of infarction and sudden death [84].

### Calcium antagonists

Calcium channel blockers, indicated in stable chronic angina, have a similar antiangina efficacy as beta-blockers [85]. The use of the rate-limiting calcium antagonists (diltiazem, verapamil) in older adults is contraindicated in the case of decompensated heart failure, severe conduction disorders and beta-blocker administration.

### Long-acting nitrates

Long-acting nitrates reduce the frequency and severity of angina attacks. No placebo-controlled study has been performed showing that nitrates have a benefit on morbidity and mortality. They reduce the frequency and severity of angina attacks, but in older adults they are also one of the main therapeutic classes responsible for orthostatic hypotension. They may be used in the case of contraindication or poor tolerability of beta-blockers and calcium antagonists [82]. Tight aortic stenosis and obstructive cardiomyopathy are contraindications to the use of nitrates.

### Other antiangina agents: molsidomine, nicorandil, trimetazidine, ivabradine

Molsidomine is a nitric-oxide donor with similar effects to nitrates. Nicorandil is a potassium channel activator with a vasodilator effect with similar effects to nitrates. The IONA [86] study showed a benefit for hospital readmissions for angina, but did not include very elderly subjects. These

two products may be used as an alternative to nitrates when these are poorly tolerated.

Trimetazidine is a metabolic antiangina drug that inhibits the beta-oxidation of fatty acids by mitochondria. It has not been shown to have an effect on prognosis [87].

The role of new heart-rate-lowering antiangina agents that inhibit the I(f) current, such as ivabradine, is not currently codified in very elderly subjects as no specific study has been performed in this population. Symptomatic treatment must be reassessed regularly and may be reduced if angina attacks become rare.

## Secondary prevention

As in younger patients, the secondary prevention of coronary events in older adults is based on the use of antithrombotic agents, beta-blockers, ACE inhibitors/ARAs and statins, and on the control of blood pressure, in combination with diet and lifestyle measures.

## Antiplatelet therapy

Treatment with platelet-aggregation inhibitors is indicated in older adults for the secondary prevention of myocardial infarction. Treatment must be continued indefinitely in the absence of haemorrhagic complications. The inhibition of platelet aggregation by aspirin effectively prevents the recurrence of myocardial infarction and cardiovascular mortality. The Antiplatelet Trialists Collaboration [88] meta-analysis showed a 27% reduction in cardiovascular complications (infarction, stroke and death from vascular causes) with aspirin. This benefit was observed irrespective of age group (> or < 65 years). Clopidogrel has proven efficacy for the prophylaxis of myocardial infarction in the over 65 subgroup of patients at high cardiovascular risk [89].

## Beta-blockers, ACE inhibitors and ARAs

The dosages of beta-blockers and inhibitors of the renin-angiotensin system must be adjusted according to their safety, and treatment continued indefinitely. Monitoring procedures are the same as during the acute phase.

Beta-blockers are recommended for the secondary prevention of myocardial infarction. Their administration reduces the risk of death in older adults who have survived myocardial infarction [90]. ACE inhibitors are also recommended for the secondary prevention of myocardial infarction in the elderly provided that they are well tolerated (in particular with respect to blood pressure and renal parameters). They are effective after myocardial infarction, with [52,91,92] or without left ventricular dysfunction [93–96] to reduce the risk of mortality and cardiovascular events. This benefit is similar whatever the age (< or > 65 years [82,85]), although these studies did not include very elderly patients.

ARAs had a similar benefit to that observed with ACE inhibitors in post-infarction complicated by heart failure or left ventricular dysfunction and in patients with coronary disease [55,97]. They may therefore be used in patients intolerant to ACE inhibitors.

## Statins

Statins are recommended for the secondary prevention of myocardial infarction in older adults. They effectively prevent coronary events in elderly subjects at high cardiovascular risk. The PROSPER study [98], which included patients aged 70 to 82 years, showed that pravastatin significantly reduced the risk of death from coronary artery disease. In the Heart Protection Study [99], which included more than 5000 patients over 75 years of age with arterial disease, cerebrovascular disease or diabetes, simvastatin reduced the incidence of a first non-fatal episode of myocardial infarction or death from coronary disease. The risk reduction was similar whatever the age. Lastly, a recent meta-analysis showed a 22% reduction in 5-year mortality in patients over 65 years receiving statins for secondary prevention [100].

## Antihypertensive medication

Antihypertensive medications effectively reduce the incidence of myocardial infarction in hypertensive subjects aged 60 to 80 years, with a risk reduction of 20–30% [101]. In patients aged over 80 years, the HYVET study showed a benefit of treatment with thiazide diuretic (indapamide) with or without an ACE inhibitor, with a 34% reduction in cardiovascular events [102]. In this study, the 28% reduction in myocardial infarction observed with treatment was found to be non-significant, possibly because of the lack of statistical power after premature discontinuation of the study.

## Lifestyle changes

Smoking cessation has a positive impact on the outcome of coronary artery disease, including in elderly subjects. Smoking cessation after myocardial infarction reduces mortality by 25–50% [103], regardless of age. Smoking cessation is possible in older adults and is achieved at rates equivalent to or higher than those observed in younger subjects.

## Physical exercise and rehabilitation

In the general population, controlled trials have demonstrated the benefit of rehabilitation in terms of functional capacities, mortality [104], reduction in atherosclerosis lesions [105] and in the number of hospital admissions for a coronary event [106]. The impact of rehabilitation in older adults was not specifically evaluated.

Rehabilitation also leads to an improvement in performance of the ADL [107]. Besides aerobic rehabilitation, resistance-training techniques (anaerobic) are effective in the elderly [108–110]. Exercises performed at home are as well tolerated and as effective as exercises performed in rehabilitation centres with medical monitoring [111].

## Management of excess body weight

Obesity is a traditional risk factor for coronary artery disease, including in the elderly [112,113]. Body mass index should be calculated in all coronary patients [114]. However, in the case of malnutrition, body mass index [115] cannot be used to recognize sarcopenic obesity (obesity with



weight loss due to loss of muscle mass), which is particularly frequent in older subjects.

As no interventional study has yet been performed, the dietetic procedures required to reduce cardiovascular risk in overweight older adults do not form the subject of a consensus. In overweight, elderly coronary patients a small reduction (500 to 700 kcal/j) in energy intake may be proposed while maintaining a daily protein intake of 1 g/kg per day in order to achieve a weight loss of 8–10% in six months [116]. Conversely, too great a weight loss in an elderly patient may have a negative effect by decreasing the muscle mass and protein reserve. Regular physical activity may increase weight loss without leading to muscle loss and should be widely recommended in elderly subjects.

### Myocardial revascularization

The type of myocardial revascularization (angioplasty or surgery) performed in older adults depends on clinical criteria (comorbidities, clinical severity) and paraclinical parameters (documented myocardial ischaemia, left ventricular function, angiographic appearance of coronary lesions). A case-by-case evaluation is necessary in these patients in order to evaluate the benefit-risk ratio.

### Coronary angioplasty

Angioplasty allows rapid treatment of coronary lesions compared with surgery, which generally requires extracorporeal circulation. Long hospital stays generating complications specific to this type of patient are therefore avoided. However, in older adults the coronary arteries needing revascularization are often the site of particularly complex lesions (extensive, diffuse, calcified lesions with frequent stenosis of the left main trunk and chronic coronary occlusions) [117].

Primary success rates vary between 80 and 96%, but revascularization is generally incomplete in patients aged over 75 years [118], with a history of infarction [119], with calcified total or sub-total coronary occlusion [120], or in the case of involvement of the circumflex artery [121].

The technical difficulties of PTCA place elderly patients in a high-risk group for per- and post-procedure complications [122–125].

Although the in-hospital complication rates and mortality are low, they are slightly higher than those of a younger population (2–3%). After 65 years of age, in-hospital mortality is independently associated with a left ventricular ejection fraction less than 40%, the existence of triple-vessel lesions and with female gender [126]. Medium-term mortality after successful PTCA is not increased in older adults

[127]. Compared with the medical strategy, PTCA has no benefit on mortality or on the recurrence of major cardiovascular events, but improves quality of life [128,129].

The indications for stenting in older adults are the same as in younger patients. The benefit compared with PTCA alone is not documented in patients aged over 80 years. The risk of bleeding induced by dual antiplatelet therapy should be carefully assessed before implanting a drug-eluting stent.

### Coronary artery bypass grafting

The benefit of surgery on survival has been demonstrated in older adults but few studies have included patients aged over 80 years [130]. Coronary artery bypass grafting has a high risk of complications (stroke, cognitive impairment, bleeding, respiratory decompensation), of approximately 30% in octogenarians [78,131].

The main predictive factors of mortality are an elderly age, female gender, resting angina, diabetes, left ventricular ejection fraction less than 40%, heart failure, renal insufficiency and emergency surgery [132]. In this setting, the appropriateness of surgical revascularization should be carefully considered by taking into account in particular the risk of deterioration of cognitive function observed after a surgical procedure.

The use of scores, such as EuroSCORE [133,134], may help to evaluate the risk of postoperative mortality, taking into account patient-related parameters (age, comorbidities), cardiopathy (left ventricular ejection fraction, unstable angina) and the surgical procedure (emergency surgery, concomitant non-coronary surgery, surgery of the chest aorta, septal rupture).

### Acknowledgements

The Editorial Committee would like to thank the company Servier, represented by Mr Pierre Schiavi and Mrs Graziella Jolly, for their help with the references and logistics. The authors wrote this document in complete scientific independence and received no financial assistance.

### Appendix 1. Mini Mental State Examination (MMSE)

Reference: Derouesne C, Poitreneau J, Hugonot L, Kalafat M, Dubois B, Laurent B. Mini-Mental State Examination: a useful method for the evaluation of the cognitive status of patients by the clinician. Consensual French version. *Presse Med* 1999;28:1141–8.

**Orientation**

*I am going to ask you a few questions to assess how well your memory works. Some are very simple, others are less so. Please answer as best you can.*

- Score/5 ☐
1. What is the year? ..... ☐
  2. What is the season? ..... ☐
  3. What is the month? ..... ☐
  4. What is the day of the month(date)?..... ☐
  5. What is the day of the week? ..... ☐

*I am now going to ask you a few questions about the place where we are at the moment:*

- Score/5 ☐
6. What is the name of the hospital we are in(if the examination is conducted at the office, asking the name of the street)? ..... ☐
  7. In which town is it? ..... ☐
  8. In which *department* (county) is this town located? ..... ☐
  9. In which region is this *department*? ..... ☐
  10. What floor are we on? ..... ☐

**Learning**

*I am going to say 3 words. I want you to repeat them to me and try and remember them because I will ask you again later.*

- Score/3 ☐
11. Cigar ..... ☐
  12. Flower ..... ☐
  13. Door ..... ☐
- Repeat the 3 words.

**Attention and calculation**

*Count backwards from 100 by subtracting 7 each time*

- Score/5 ☐
14. 93 ..... ☐
  15. 86 ..... ☐
  16. 79 ..... ☐
  17. 72 ..... ☐
  18. 65 ..... ☐

For all subjects, even those with the maximum points, ask: Please spell the word *MONDE* backwards: EDNOM.

The score is the number of letters in the right place. (This figure must not appear in the global score.)

**Recall**

Can you tell me the 3 words that I asked you to repeat and remember a few moments ago?


Score/3 ☐

19. Cigar ..... ☐  
 20. Flower ..... ☐  
 21. Door ..... ☐

**Language**

Score/9 ☐

22. Show the patient a pencil. *What is this called?* ..... ☐  
 23. Show your watch. *What is this called?* ..... ☐  
 24. *Listen carefully and repeat after me: "No ifs, ands or buts"* ..... ☐  
 25. Place a sheet of paper on the desk and show it to the subject while saying: *Listen carefully and do what I tell you to do:*  
     Take this paper sheet with the right hand, ..... ☐  
 26. Fold it in half ..... ☐  
 27. And throw it on the floor ..... ☐  
 28. Give the subject a sheet of paper with the following written in large letters: **"Close your eyes"**  
     and tell the subject:  
     *Do what it says* ..... ☐  
 29. Give the subject a sheet of paper and a pen and tell the patient: *Write any sentence that you want though it must be a whole sentence. This sentence must be written spontaneously. It must contain a subject, a verb and make sense.* ☐  
**Copying**  
 30. Give the subject a sheet of paper and ask him or her:  
     *"Can you copy this drawing?"* ☐



Score total (0 to 30) :  
 NORMAL SCORE =  $25 \pm 2$

## Appendix 2. Simplified IADL Scale (PAQUID Study)

Reference: Barberger-Gateau P, Dartigues JF, Letenneur L. Four Instrumental Activities of Daily Living Score as a predictor of one-year incident dementia. *Age Ageing* 1993;22:457–63.

### 1. Ability to use the telephone

- ☐ Makes use normally of the telephone (telephones on his/her own initiative, composes numbers)

### 2. Means of transport

- ☐ Uses public transport independently or drives his/her own car

### 3. Responsibility for own medications

- ☐ Deals with his/her own medications personally (correct doses and schedules)

### 4. Ability to handle finances

- ☐ Manages his or her budget autonomously (writes cheques, pays the rent and bills, bank operations)

Each of the 4 items scored (0) = need for help, (1) = no need for help, according to the response.

Score/4 ☐

(normal score = 4/4)



### Appendix 3. Katz ADL Scale

Reference: Katz A, Ford AB, Moskowitz RW et al. *Studies of illness in the aged. The index of ADL: a standardized measure of biological and psychosocial function.* J Am Med Assoc 1963;185:914–9

Score  
If total score equal or greater than 1, strong probability of depression  
If total score = 0, strong probability of no depression

#### Bathing (wash-basin, bath or shower and body hygiene)

- 1 ☐ No need for help  
½ ☐ Partial need for help  
0 ☐ Dependence

#### Dressing (gets clothes from cupboards or drawers, including underwear and outer garments and uses button and zip fasteners)

- 1 ☐ No need for help  
½ ☐ Autonomous for the choice of clothes and getting dressed but may need help tying shoes  
0 ☐ Dependence

#### Toileting (to urinate or defecate, wipe themselves and get dressed)

- 1 ☐ No need for help  
½ ☐ Must be accompanied or needs help to undress and put clothes back on  
0 ☐ Can't go to the toilets alone or doesn't use the bedpan

#### Transferring

- 1 ☐ Moves in and out of bed or chair unassisted (may use a stick or walker)  
½ ☐ Needs help  
0 ☐ Stays in bed (bedridden)

#### Continence

- 1 ☐ Complete self control over urination and defecation  
½ ☐ Occasional accidents  
0 ☐ Completely incontinent

#### Feeding

- 1 ☐ No need for help  
½ ☐ Needs help cutting meat, buttering bread or peeling fruit  
0 ☐ Needs total help with feeding

Score/6 ☐

(Normal score = 6/6)

### Appendix 4. Mini GDS (Geriatric Depression Scale), mood assessment

Reference: Clément J-P, Nassif RF, Leger JM, Marchan F. Mise au point et contribution à la validation d'une version française brève de la Geriatric Depression Scale de Yesavage. L'Encéphale 1997;XXIII:91–9.

Ask the patient to answer the questions by saying how he/she has felt during the last week and not during his life or at the present time.

- |   |                 |
|---|-----------------|
| 1. Do you often feel discouraged and sad?                       | yes = 1, no = 0 |
| 2. Do you feel that your life is empty?                         | yes = 1, no = 0 |
| 3. Are you happy most of the time?                              | yes = 0, no = 1 |
| 4. Do you have the impression that your situation is desperate? | yes = 1, no = 0 |

### References

- [1] National Federation of Regional Public Health Observatories. Cardiovascular disease. [www.fnors.org/Fnors/Ors/Travaux/So/703.pdf](http://www.fnors.org/Fnors/Ors/Travaux/So/703.pdf).
- [2] Alexander KP, Roe MT, Chen AY, et al. Evolution in cardiovascular care for elderly patients with non-ST-segment elevation acute coronary syndromes: results from the CRUSADE National Quality Improvement Initiative. J Am Coll Cardiol 2005;46:1479–87.
- [3] Avezum A, Makdisse M, Spencer F, et al. Impact of age on management and outcome of acute coronary syndrome: observations from the Global Registry of Acute Coronary Events (GRACE). Am Heart J 2005;149:67–73.
- [4] Goldberg RJ, Yarzebski J, Lessard D, et al. A two-decades (1975 to 1995) long experience in the incidence, in-hospital and long-term case-fatality rates of acute myocardial infarction: a community-wide perspective. J Am Coll Cardiol 1999;33:1533–9.

- [5] Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction-executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to revise the 1999 guidelines for the management of patients with acute myocardial infarction). *J Am Coll Cardiol* 2004;44:671-719.
- [6] Bertrand ME, Simoons ML, Fox KA, et al. Management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. The Task Force on the management of acute coronary syndromes of the European Society of Cardiology. *Eur Heart J* 2002;23:1809-40.
- [7] Aronow WS, Ahn C, Mercando AD, et al. Prevalence of and association between silent myocardial ischemia and new coronary events in older men and women with and without cardiovascular disease. *J Am Geriatr Soc* 2002;50:1075-8.
- [8] Cohn PF, Fox KM, Daly C. Silent myocardial ischemia. *Circulation* 2003;108:1263-77.
- [9] Gill TM, DiPietro L, Krumholz HM. Role of exercise stress testing and safety monitoring for older persons starting an exercise program. *JAMA* 2000;284:342-9.
- [10] Amendo MT, Brown BA, Kossow LB, et al. Headache as the sole presentation of acute myocardial infarction in two elderly patients. *Am J Geriatr Cardiol* 2001;10:100-1.
- [11] Dangelser G, Gottwalles Y, Huk M, et al. Acute ST-elevation myocardial infarction in the elderly (> 75 years). Results from a regional multicenter study. *Presse Med* 2005;34:983-9.
- [12] Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189-98.
- [13] Derouesne C, Poitreneau J, Hugonot L, et al. Mini-Mental State Examination: a useful method for the evaluation of the cognitive status of patients by the clinician. Consensual French version. *Presse Med* 1999;28:1141-8.
- [14] Krumholz HM, Chen J, Chen YT, et al. Predicting one-year mortality among elderly survivors of hospitalization for an acute myocardial infarction: results from the Cooperative Cardiovascular Project. *J Am Coll Cardiol* 2001;38:453-9.
- [15] Barberger-Gateau P, Dartigues JF, Letenneur L. Four Instrumental Activities of Daily Living Score as a predictor of one-year incident dementia. *Age Ageing* 1993;22:457-63.
- [16] Katz S, Ford AB, Moskowitz RW, et al. Studies of Illness in the Aged. The Index of Adl: A Standardized Measure of Biological and Psychosocial Function. *JAMA* 1963;185:914-9.
- [17] Guigoz Y, Lauque S, Vellas BJ. Identifying the elderly at risk for malnutrition. The Mini Nutritional Assessment. *Clin Geriatr Med* 2002;18:737-57.
- [18] Ariyo AA, Haan M, Tangen CM, et al. Depressive symptoms and risks of coronary heart disease and mortality in elderly Americans. *Circulation* 2000;102:1773-9.
- [19] Rubenstein LZ, Josephson KR, Wieland GD, et al. Effectiveness of a geriatric evaluation unit. A randomized clinical trial. *N Engl J Med* 1984;311:1664-70.
- [20] Gibbons RJ, Chatterjee K, Daley J, et al. ACC/AHA/ACP-ASIM guidelines for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Patients With Chronic Stable Angina). *J Am Coll Cardiol* 1999;33:2092-197.
- [21] Aronow WS. Correlation of ischemic ST-segment depression on the resting electrocardiogram with new cardiac events in 1,106 patients over 62 years of age. *Am J Cardiol* 1989;64:232-3.
- [22] Hollenberg M, Ngo LH, Turner D, et al. Treadmill exercise testing in an epidemiologic study of elderly subjects. *J Gerontol A Biol Sci Med Sci* 1998;53:B259-67.
- [23] Kwok JM, Miller TD, Hodge DO, et al. Prognostic value of the Duke treadmill score in the elderly. *J Am Coll Cardiol* 2002;39:1475-81.
- [24] Jeger RV, Zellweger MJ, Kaiser C, et al. Prognostic value of stress testing in patients over 75 years of age with chronic angina. *Chest* 2004;125:1124-31.
- [25] Wang FP, Amanullah AM, Kiat H, et al. Diagnostic efficacy of stress technetium 99m-labeled sestamibi myocardial perfusion single-photon emission computed tomography in detection of coronary artery disease among patients over age 80. *J Nucl Cardiol* 1995;2:380-8.
- [26] Solodky A, Assali AR, Mats I, et al. Prognostic value of myocardial perfusion imaging in symptomatic and asymptomatic patients after percutaneous coronary intervention. *Cardiology* 2007;107:38-43.
- [27] Dhond MR, Nguyen TT, Sabapathy R, et al. Dobutamine stress echocardiography in preoperative and long-term postoperative risk assessment of elderly patients. *Am J Geriatr Cardiol* 2003;12:107-9, 12.
- [28] Hiro J, Hiro T, Reid CL, et al. Safety and results of dobutamine stress echocardiography in women versus men and in patients older and younger than 75 years of age. *Am J Cardiol* 1997;80:1014-20.
- [29] Serdar MA, Tokgoz S, Metinyurt G, et al. Effect of macro-creatine kinase and increased creatine kinase BB on the rapid diagnosis of patients with suspected acute myocardial infarction in the emergency department. *Mil Med* 2005;170:648-52.
- [30] Babuin L, Jaffe AS. Troponin: the biomarker of choice for the detection of cardiac injury. *CMAJ* 2005;173:1191-202.
- [31] Thygesen K, Alpert JS, White HD. Universal definition of myocardial infarction. *Eur Heart J* 2007;28:2525-38.
- [32] Alexander KP, Newby LK, Armstrong PW, et al. Acute coronary care in the elderly, part II: ST-segment-elevation myocardial infarction: a scientific statement for healthcare professionals from the American Heart Association Council on Clinical Cardiology: in collaboration with the Society of Geriatric Cardiology. *Circulation* 2007;115:2570-89.
- [33] Alexander KP, Newby LK, Cannon CP, et al. Acute coronary care in the elderly, part I: Non-ST-segment-elevation acute coronary syndromes: a scientific statement for healthcare professionals from the American Heart Association Council on Clinical Cardiology: in collaboration with the Society of Geriatric Cardiology. *Circulation* 2007;115:2549-69.
- [34] Bassand JP, Hamm CW, Ardissino D, et al. Guidelines for the diagnosis and treatment of non-ST-segment elevation acute coronary syndromes. *Eur Heart J* 2007;28:1598-660.
- [35] Braunwald E, Antman EM, Beasley JW, et al. ACC/AHA guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction-2002: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on the Management of Patients With Unstable Angina). *Circulation* 2002;106:1893-900.
- [36] Van de Werf F, Bax J, Betriu A, et al. Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: The Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology. *Eur Heart J* 2008;29:2909-45.
- [37] Yusuf S, Zhao F, Mehta SR, et al. Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med* 2001;345:494-502.
- [38] Mehta SR, Yusuf S, Peters RJ, et al. Effects of pretreatment with clopidogrel and aspirin followed by long-term therapy in patients undergoing percutaneous coronary intervention: the PCI-CURE study. *Lancet* 2001;358:527-33.

- [39] Albaladejo P, Marret E, Piriou V, et al. Management of oral antiplatelet treatment for patients with coronary stents. *Ann Fr Anesth Reanim* 2006;25:796–8.
- [40] Montalescot G, Barragan P, Wittenberg O, et al. Platelet glycoprotein IIb/IIIa inhibition with coronary stenting for acute myocardial infarction. *N Engl J Med* 2001;344:1895–903.
- [41] Bach RG, Cannon CP, Weintraub WS, et al. The effect of routine, early invasive management on outcome for elderly patients with non-ST-segment elevation acute coronary syndromes. *Ann Intern Med* 2004;141:186–95.
- [42] Wallentin L, Goldstein P, Armstrong PW, et al. Efficacy and safety of tenecteplase in combination with the low-molecular-weight heparin enoxaparin or unfractionated heparin in the prehospital setting: the Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)-3 PLUS randomized trial in acute myocardial infarction. *Circulation* 2003;108:135–42.
- [43] Murphy SA, Gibson CM, Morrow DA, et al. Efficacy and safety of the low-molecular weight heparin enoxaparin compared with unfractionated heparin across the acute coronary syndrome spectrum: a meta-analysis. *Eur Heart J* 2007;28:2077–86.
- [44] Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)-3 Investigators. Efficacy and safety of tenecteplase in combination with enoxaparin, abciximab, or unfractionated heparin: the ASSENT-3 randomised trial in acute myocardial infarction. *Lancet* 2001;358:605–13.
- [45] Cohen M, Demers C, Gurfinkel EP, et al. Low-molecular-weight heparins in non-ST-segment elevation ischemia: the ESSENCE trial. Efficacy and Safety of Subcutaneous Enoxaparin versus intravenous unfractionated heparin, in non-Q-wave Coronary Events. *Am J Cardiol* 1998;82:19L–24L.
- [46] Stone GW, White HD, Ohman EM, et al. Bivalirudin in patients with acute coronary syndromes undergoing percutaneous coronary intervention: a subgroup analysis from the Acute Catheterization and Urgent Intervention Triage strategy (ACUITY) trial. *Lancet* 2007;369:907–19.
- [47] Lincoff AM, Kleiman NS, Kereiakes DJ, et al. Long-term efficacy of bivalirudin and provisional glycoprotein IIb/IIIa blockade vs heparin and planned glycoprotein IIb/IIIa blockade during percutaneous coronary revascularization: REPLACE-2 randomized trial. *Jama* 2004;292:696–703.
- [48] Yusuf S, Mehta SR, Chrolavicius S, et al. Comparison of fondaparinux and enoxaparin in acute coronary syndromes. *N Engl J Med* 2006;354:1464–76.
- [49] Yusuf S, Mehta SR, Chrolavicius S, et al. Effects of fondaparinux on mortality and reinfarction in patients with acute ST-segment elevation myocardial infarction: the OASIS-6 randomized trial. *JAMA* 2006;295:1519–30.
- [50] Dargie HJ. Effect of carvedilol on outcome after myocardial infarction in patients with left-ventricular dysfunction: the CAPRICORN randomised trial. *Lancet* 2001;357:1385–90.
- [51] Flather MD, Yusuf S, Kober L, et al. Long-term ACE-inhibitor therapy in patients with heart failure or left-ventricular dysfunction: a systematic overview of data from individual patients. *Lancet* 2000;355:1575–81.
- [52] Pfeffer MA, Braunwald E, Moye LA, et al. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. Results of the survival and ventricular enlargement trial. *N Engl J Med* 1992;327:669–77.
- [53] Krumholz HM, Chen YT, Wang Y, et al. Aspirin and angiotensin-converting enzyme inhibitors among elderly survivors of hospitalization for an acute myocardial infarction. *Arch Intern Med* 2001;161:538–44.
- [54] Pfeffer MA, McMurray JJ, Velazquez EJ, et al. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003;349:1893–906.
- [55] Dickstein K, Kjeksus J. Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomised trial. Optimal Trial in Myocardial Infarction with Angiotensin II Antagonist Losartan. *Lancet* 2002;360:752–60.
- [56] Pitt B, Remme W, Zannad F, et al. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003;348:1309–21.
- [57] Ray KK, Bach RG, Cannon CP, et al. Benefits of achieving the NCEP optional LDL-C goal among elderly patients with ACS. *Eur Heart J* 2006;27:2310–6.
- [58] ISIS-4 (Fourth International Study of Infarct Survival) Collaborative Group. ISIS-4: a randomised factorial trial assessing early oral captopril, oral mononitrate, and intravenous magnesium sulphate in 58,050 patients with suspected acute myocardial infarction. *Lancet* 1995;345:669–85.
- [59] Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico. Six-month effects of early treatment with lisinopril and transdermal glyceryl trinitrate singly and together withdrawn six weeks after acute myocardial infarction: the GISSI-3 trial. *J Am Coll Cardiol* 1996;27:337–44.
- [60] The GUSTO investigators. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. *N Engl J Med* 1993;329:673–82.
- [61] Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico. GISSI-2: a factorial randomised trial of alteplase versus streptokinase and heparin versus no heparin among 12,490 patients with acute myocardial infarction. *Lancet* 1990;336:65–71.
- [62] ISIS-3 (Third International Study of Infarct Survival) Collaborative Group. ISIS-3: a randomised comparison of streptokinase vs tissue plasminogen activator vs anistreplase and of aspirin plus heparin vs aspirin alone among 41,299 cases of suspected acute myocardial infarction. *Lancet* 1992;339:753–70.
- [63] Mehta RH, Rathore SS, Radford MJ, et al. Acute myocardial infarction in the elderly: differences by age. *J Am Coll Cardiol* 2001;38:736–41.
- [64] Rogers WJ, Bowlby LJ, Chandra NC, et al. Treatment of myocardial infarction in the United States (1990 to 1993). Observations from the National Registry of Myocardial Infarction. *Circulation* 1994;90:2103–14.
- [65] Thiemann DR, Coresh J, Schulman SP, et al. Lack of benefit for intravenous thrombolysis in patients with myocardial infarction who are older than 75 years. *Circulation* 2000;101:2239–46.
- [66] Berger AK, Radford MJ, Wang Y, et al. Thrombolytic therapy in older patients. *J Am Coll Cardiol* 2000;36:366–74.
- [67] White HD, Barbash GI, Califf RM, et al. Age and outcome with contemporary thrombolytic therapy. Results from the GUSTO-I trial. Global Utilization of Streptokinase and TPA for Occluded coronary arteries trial. *Circulation* 1996;94:1826–33.
- [68] Mehta RH, Granger CB, Alexander KP, et al. Reperfusion strategies for acute myocardial infarction in the elderly: benefits and risks. *J Am Coll Cardiol* 2005;45:471–8.
- [69] Tiefenbrunn AJ, Chandra NC, French WJ, et al. Clinical experience with primary percutaneous transluminal coronary angioplasty compared with alteplase (recombinant tissue-type plasminogen activator) in patients with acute myocardial infarction: a report from the Second National Registry of Myocardial Infarction (NRM-2). *J Am Coll Cardiol* 1998;31:1240–5.
- [70] Berger AK, Schulman KA, Gersh BJ, et al. Primary coronary angioplasty vs thrombolysis for the management of acute myocardial infarction in elderly patients. *JAMA* 1999;282:341–8.
- [71] Zahn R, Schiele R, Schneider S, et al. Primary angioplasty versus intravenous thrombolysis in acute myocardial infarction.



- tion: can we define subgroups of patients benefiting most from primary angioplasty? Results from the pooled data of the Maximal Individual Therapy in Acute Myocardial Infarction Registry and the Myocardial Infarction Registry. *J Am Coll Cardiol* 2001;37:1827–35.
- [72] Mehta RH, Sadiq I, Goldberg RJ, et al. Effectiveness of primary percutaneous coronary intervention compared with that of thrombolytic therapy in elderly patients with acute myocardial infarction. *Am Heart J* 2004;147:253–9.
- [73] Batchelor WB, Anstrom KJ, Muhlbaier LH, et al. Contemporary outcome trends in the elderly undergoing percutaneous coronary interventions: results in 7,472 octogenarians. *J Am Coll Cardiol* 2000;36:723–30.
- [74] Hochman JS, Sleeper LA, Webb JG, et al. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. SHOCK Investigators. Should we emergently revascularize occluded coronaries for cardiogenic shock. *N Engl J Med* 1999;341:625–34.
- [75] Dzavik V, Sleeper LA, Cocke TP, et al. Early revascularization is associated with improved survival in elderly patients with acute myocardial infarction complicated by cardiogenic shock: a report from the SHOCK Trial Registry. *Eur Heart J* 2003;24:828–37.
- [76] Dauerman HL, Goldberg RJ, Malinski M, et al. Outcomes and early revascularization for patients > or =65 years of age with cardiogenic shock. *Am J Cardiol* 2001;87:844–8.
- [77] Dauerman HL, Ryan Jr TJ, Piper WD, et al. Outcomes of percutaneous coronary intervention among elderly patients in cardiogenic shock: a multicenter, decade-long experience. *J Invasive Cardiol* 2003;15:380–4.
- [78] Cosgrove DM. Surgical revascularization in the elderly. *Cardiol Elderly* 1993;1:71–6.
- [79] Kennedy JW, Ivey TD, Misbach G, et al. Coronary artery bypass graft surgery early after acute myocardial infarction. *Circulation* 1989;79:173–8.
- [80] Fox K, Garcia MA, Ardissino D, et al. Guidelines on the management of stable angina pectoris: executive summary: The Task Force on the Management of Stable Angina Pectoris of the European Society of Cardiology. *Eur Heart J* 2006;27:1341–81.
- [81] Abrams J. Clinical practice. Chronic stable angina. *N Engl J Med* 2005;352:2524–33.
- [82] Fihn SD, Williams SV, Daley J, et al. Guidelines for the management of patients with chronic stable angina: treatment. *Ann Intern Med* 2001;135:616–32.
- [83] Gibbons RJ, Abrams J, Chatterjee K, et al. ACC/AHA 2002 guideline update for the management of patients with chronic stable angina-summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on the Management of Patients With Chronic Stable Angina). *Circulation* 2003;107:149–58.
- [84] Pharmacotherapy BDIC. Management of stable angina. Transparency Sheet Update 2005: [www.cbip.be/PDF/TFT/TF\\_ANGR.pdf](http://www.cbip.be/PDF/TFT/TF_ANGR.pdf).
- [85] Heidenreich PA, McDonald KM, Hastie T, et al. Meta-analysis of trials comparing beta-blockers, calcium antagonists, and nitrates for stable angina. *JAMA* 1999;281:1927–36.
- [86] IONA Study Group. Effect of nicorandil on coronary events in patients with stable angina: the Impact of Nicorandil in Angina (IONA) randomised trial. *Lancet* 2002;359:1269–75.
- [87] Stanley WC, Marzilli M. Metabolic therapy in the treatment of ischaemic heart disease: the pharmacology of trimetazidine. *Fundam Clin Pharmacol* 2003;17:133–45.
- [88] Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy-I: Prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. *BMJ* 1994;308:81–106.
- [89] Cannon CP. Effectiveness of clopidogrel versus aspirin in preventing acute myocardial infarction in patients with symptomatic atherothrombosis (CAPRIE trial). *Am J Cardiol* 2002;90:760–2.
- [90] Gottlieb SS, McCarter RJ, Vogel RA. Effect of beta-blockade on mortality among high-risk and low-risk patients after myocardial infarction. *N Engl J Med* 1998;339:489–97.
- [91] The Acute Infarction Ramipril Efficacy (AIRE) Study Investigators. Effect of ramipril on mortality and morbidity of survivors of acute myocardial infarction with clinical evidence of heart failure. *Lancet* 1993;342:821–8.
- [92] Kober L, Torp-Pedersen C, Carlsen JE, et al. A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 1995;333:1670–6.
- [93] Yusuf S, Sleight P, Pogue J, et al. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med* 2000;342:145–53.
- [94] Fox KM. Efficacy of perindopril in reduction of cardiovascular events among patients with stable coronary artery disease: randomised, double-blind, placebo-controlled, multicentre trial (the EUROPA study). *Lancet* 2003;362:782–8.
- [95] Dagenais GR, Pogue J, Fox K, et al. Angiotensin-converting-enzyme inhibitors in stable vascular disease without left ventricular systolic dysfunction or heart failure: a combined analysis of three trials. *Lancet* 2006;368:581–8.
- [96] Danchin N, Cucherat M, Thillez C, et al. Angiotensin-converting enzyme inhibitors in patients with coronary artery disease and absence of heart failure or left ventricular systolic dysfunction: an overview of long-term randomized controlled trials. *Arch Intern Med* 2006;166:787–96.
- [97] Yusuf S, Teo KK, Pogue J, et al. Telmisartan, ramipril, or both in patients at high risk for vascular events. *N Engl J Med* 2008;358:1547–59.
- [98] Shepherd J, Blauw GJ, Murphy MB, et al. Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *Lancet* 2002;360:1623–30.
- [99] Collins R, Armitage J, Parish S, et al. Effects of cholesterol-lowering with simvastatin on stroke and other major vascular events in 20536 people with cerebrovascular disease or other high-risk conditions. *Lancet* 2004;363:757–67.
- [100] Afalo J, Duque G, Steele R, et al. Statins for secondary prevention in elderly patients: a hierarchical bayesian meta-analysis. *J Am Coll Cardiol* 2008;51:37–45.
- [101] Insua JT, Sacks HS, Lau TS, et al. Drug treatment of hypertension in the elderly: a meta-analysis. *Ann Intern Med* 1994;121:355–62.
- [102] Beckett NS, Peters R, Fletcher AE, et al. Treatment of hypertension in patients 80 years of age or older. *N Engl J Med* 2008;358:1887–98.
- [103] Sparrow D, Dawber TR. The influence of cigarette smoking on prognosis after a first myocardial infarction. A report from the Framingham study. *J Chronic Dis* 1978;31:425–32.
- [104] O'Connor GT, Buring JE, Yusuf S, et al. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. *Circulation* 1989;80:234–44.
- [105] Haskell WL, Alderman EL, Fair JM, et al. Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP). *Circulation* 1994;89:975–90.
- [106] Ornish D, Scherwitz LW, Billings JH, et al. Intensive lifestyle changes for reversal of coronary heart disease. *JAMA* 1998;280:2001–7.
- [107] Ades PA, Maloney A, Savage P, et al. Determinants of physical functioning in coronary patients: response to cardiac rehabilitation. *Arch Intern Med* 1999;159:2357–60.



- [108] Pollock ML, Franklin BA, Balady GJ, et al. AHA Science Advisory. Resistance exercise in individuals with and without cardiovascular disease: benefits, rationale, safety, and prescription: An advisory from the Committee on Exercise, Rehabilitation, and Prevention, Council on Clinical Cardiology, American Heart Association; Position paper endorsed by the American College of Sports Medicine. *Circulation* 2000;101:828–33.
- [109] American College of Sports Medicine Position Stand. Exercise for patients with coronary artery disease. *Med Sci Sports Exerc* 1994;26:i–v.
- [110] American Association of Cardiovascular and Pulmonary Rehabilitation. Guidelines for Cardiac Rehabilitation and Secondary Programs. 4th ed. Champaign Ill ed Human kinetics; 2004.
- [111] Harris DE, Record NB. Cardiac rehabilitation in community settings. *J Cardiopulm Rehabil* 2003;23:250–9.
- [112] Song Y, Manson JE, Meigs JB, et al. Comparison of usefulness of body mass index versus metabolic risk factors in predicting 10-year risk of cardiovascular events in women. *Am J Cardiol* 2007;100:1654–8.
- [113] He Y, Jiang B, Wang J, et al. BMI versus the metabolic syndrome in relation to cardiovascular risk in elderly Chinese individuals. *Diabetes Care* 2007;30:2128–34.
- [114] Montaye M, De Bacquer D, De Backer G, et al. Overweight and obesity: a major challenge for coronary heart disease secondary prevention in clinical practice in Europe. *Eur Heart J* 2000;21:808–13.
- [115] Zamboni M, Mazzali G, Zoico E, et al. Health consequences of obesity in the elderly: a review of four unresolved questions. *Int J Obes (Lond)* 2005;29:1011–29.
- [116] Villareal DT, Apovian CM, Kushner RF, et al. Obesity in older adults: technical review and position statement of the American Society for Nutrition and NAAO. The Obesity Society. *Am J Clin Nutr* 2005;82:923–34.
- [117] Dynina O, Vakili BA, Slater JN, et al. In-hospital outcomes of contemporary percutaneous coronary interventions in the very elderly. *Catheter Cardiovasc Interv* 2003;58:351–7.
- [118] Klein LW, Block P, Brindis RG, et al. Percutaneous coronary interventions in octogenarians in the American College of Cardiology-National Cardiovascular Data Registry: development of a nomogram predictive of in-hospital mortality. *J Am Coll Cardiol* 2002;40:394–402.
- [119] Smith Jr SC, Feldman TE, Hirshfeld Jr JW, et al. ACC/AHA/SCAI 2005 guideline update for percutaneous coronary intervention: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/SCAI Writing Committee to Update the 2001 Guidelines for Percutaneous Coronary Intervention). *J Am Coll Cardiol* 2006;47:e1–121.
- [120] Chauhan MS, Kuntz RE, Ho KL, et al. Coronary artery stenting in the aged. *J Am Coll Cardiol* 2001;37:856–62.
- [121] Kaiser C, Kuster GM, Erne P, et al. Risks and benefits of optimised medical and revascularisation therapy in elderly patients with angina-on-treatment analysis of the TIME trial. *Eur Heart J* 2004;25:1036–42.
- [122] Graham MM, Ghali WA, Faris PD, et al. Survival after coronary revascularization in the elderly. *Circulation* 2002;105:2378–84.
- [123] Thompson RC, Holmes Jr DR, Gersh BJ, et al. Percutaneous transluminal coronary angioplasty in the elderly: early and long-term results. *J Am Coll Cardiol* 1991;17:1245–50.
- [124] Buffet P, Selton-Suty C, Juilliere Y, et al. Coronary transluminal angioplasty after 70 years of age. Multivariate analysis of parameters influencing immediate results and long-term prognosis. *Arch Mal Coeur Vaiss* 1992;85:287–93.
- [125] Boden WE, O'Rourke RA, Teo KK, et al. Optimal medical therapy with or without PCI for stable coronary disease. *N Engl J Med* 2007;356:1503–16.
- [126] Imburgia M, King TR, Soffer AD, et al. Early results and long-term outcome of percutaneous transluminal coronary angioplasty in patients age 75 years or older. *Am J Cardiol* 1989;63:1127–9.
- [127] Little T, Milner MR, Lee K, et al. Late outcome and quality of life following percutaneous transluminal coronary angioplasty in octogenarians. *Catheter Cardiovasc Diagn* 1993;29:261–6.
- [128] The TIME Investigators. Trial of invasive versus medical therapy in elderly patients with chronic symptomatic coronary-artery disease (TIME): a randomised trial. *Lancet* 2001;358:951–7.
- [129] Pfisterer M, Buser P, Osswald S, et al. Outcome of elderly patients with chronic symptomatic coronary artery disease with an invasive vs optimized medical treatment strategy: one-year results of the randomized TIME trial. *Jama* 2003;289:1117–23.
- [130] Filsofi F, Rahmanian PB, Castillo JG, et al. Results and predictors of early and late outcomes of coronary artery bypass graft surgery in octogenarians. *J Cardiothorac Vasc Anesth* 2007;21:784–92.
- [131] Jones EL. Coronary bypass surgery in the elderly. In: Wenger NK, Furberg CD, Pitt E, editors. *Coronary heart disease in the elderly*. New York: Elsevier Science Publishing Co., Inc; 1986. p. 375–88.
- [132] Peterson ED, Cowper PA, Jollis JG, et al. Outcomes of coronary artery bypass graft surgery in 24,461 patients aged 80 years or older. *Circulation* 1995;92:II85–91.
- [133] Nashef SA, Roques F, Michel P, et al. European system for cardiac operative risk evaluation (EuroSCORE). *Eur J Cardiothorac Surg* 1999;16:9–13.
- [134] Gogbashian A, Sedrakyan A, Treasure T. EuroSCORE: a systematic review of international performance. *Eur J Cardiothorac Surg* 2004;25:695–700.